

*Title:*

## **Relation Between the BDMPs Transport Coefficient and the Dipole Cross Section**

*Author(s):*

**Jörg Raufeisen**

*Submitted to:*

<http://lib-www.lanl.gov/cgi-bin/getfile?00818824.pdf>

# RELATION BETWEEN THE BDMPS TRANSPORT COEFFICIENT AND THE DIPOLE CROSS SECTION

Jörg Raufeisen

Los Alamos National Laboratory, MS H846, Los Alamos, NM 87545, jorgr@lanl.gov

## Abstract

In the BDMPS formalism, the transverse momentum accumulated by a quark propagating through nuclear matter is proportional to the so-called transport coefficient  $\hat{q}$ . On the other hand, transverse momentum broadening can also be calculated within the color dipole approach where it is expressed in terms of the dipole cross section  $\sigma_{q\bar{q}}$ . Since both approaches are equivalent, it is possible to find a relation between  $\hat{q}$  and  $\sigma_{q\bar{q}}$ .

Transverse momentum broadening of a fast parton (*i.e.* quark or gluon) propagating through nuclear matter has received much attention during the past years, because it is intimately related to the medium induced energy loss of that parton [1, 2]. Indeed, it was found by the BDMPS collaboration [1] that in QCD the induced radiative energy loss per unit length is proportional to the transverse momentum acquired by the parton. This leads to the seemingly counterintuitive result  $dE/dx \propto L$ , where  $L$  is the length of the medium traversed by the parton. The same result was obtained by Zakharov [3] in the color dipole approach. Even though these two approaches use very different languages to describe induced energy loss, it was found in [4] that they are equivalent. In this note, we present a quantitative relation between the two main non-perturbative inputs of both approaches, the BDMPS transport coefficient  $\hat{q}$  and the dipole cross section  $\sigma_{q\bar{q}}$ , Eq. (5). To our knowledge, this relation was first published in [5].

Broadening of transverse momentum of a fast quark propagating in nuclear matter (but not energy loss) was first investigated within the dipole approach in [6]. The authors of [6] study broadening of Drell-Yan pairs in hadron-nucleus collisions. The result of this phenomenological analysis is that the projectile quark performs a random walk as it propagates through the nucleus and thereby acquires the transverse momentum

$$\delta\langle p_T^2 \rangle = C(\tilde{x}, \tilde{Q}^2) \rho_A L, \quad (1)$$

where  $\rho_A$  is the nuclear density. The factor  $C(\tilde{x}, \tilde{Q}^2)$  originates from the dipole cross section, which can be written for small separations  $\rho$  as

$$\sigma_{q\bar{q}}(\tilde{x}, \tilde{Q}^2) = C(\tilde{x}, \tilde{Q}^2) \rho^2. \quad (2)$$

A theoretically more profound derivation of Eq. (1) can be found in [5, 7]. Note that  $C$  depends on energy  $\tilde{x}$  and on a hard scale  $\tilde{Q}^2$ . At leading order, these scales cannot be calculated exactly and one has to rely on plausible arguments to find their values, see [7]. The coefficient  $C$  can be expressed in terms of the gluon density  $G_N$  of a nucleon [8],

$$C(\tilde{x}, \tilde{Q}^2) = \frac{\pi^2}{3} \alpha_s(\lambda/\rho^2) \tilde{x} G_N(\tilde{x}/\tilde{Q}^2), \quad (3)$$

where  $\lambda$  is a dimensionless number. Note that in order to obtain broadening for gluons, one would have to multiply  $C$  by the ratio of Casimir factors for adjoint and fundamental representation of  $\mathcal{G}$  (3), *i.e.*  $C_A/C_F = 9/4$ .

On the other hand, for cold nuclear matter the BDMPS transport coefficient  $\hat{q}$  can also be expressed in terms of the gluon density of a nucleon [1]. For the quark case we are interested in, one finds

$$\hat{q} = \frac{2\pi^2 \alpha_s(Q'^2)}{3} \rho_A x' G_N(x', Q'^2). \quad (4)$$

Again, the scales  $x'$  and  $Q'^2$  cannot be determined exactly. The transport coefficient describes the “scattering power” of the medium. Our final result is obtained by comparing Eqs. (3) and (4), which yields

$$\hat{q}(x', Q'^2) = 2\rho_A C(\tilde{x}, \tilde{Q}^2). \quad (5)$$

This equality holds to logarithmic accuracy because of the scale dependence of  $\hat{q}$  and  $C$ . In addition, one finds from Eqs. (1), (3) and (4)

$$\delta\langle p_T^2 \rangle = \hat{q}L = p_{\perp W}^2, \quad (6)$$

where (following the notation of [1])  $p_{\perp W}^2$  is the characteristic transverse momentum squared of a parton produced in the medium after traversing nuclear matter of length  $L$ . The second equality in Eq. (6) was found within the BDMPS formalism in [1].

We stress that Eq. (5) was not obtained by equating Eq. (1) with the corresponding result from the BDMPS approach, *i.e.*  $p_{\perp W}^2 = \hat{q}L$ . Instead, we have shown that  $\delta\langle p_T^2 \rangle = p_{\perp W}^2$ , confirming the equivalence of the two approaches. Note that it is not clear a priori that these two quantities should be equal, since  $\delta\langle p_T^2 \rangle$  is the broadening of an incident quark, while  $p_{\perp W}^2$  is the broadening for a quark produced in the medium. However, since both quantities are proportional to the length of the medium, the result  $\delta\langle p_T^2 \rangle = p_{\perp W}^2$  is plausible.

As a concluding remark, we mention that one could attempt to set  $\delta\langle p_T^2 \rangle$  equal to the nuclear broadening obtained in the higher twist factorization approach of [9]. However, even though the relation between BDMPS/color dipole formalism and LQS approach is not fully understood yet, the LQS approach probably contains different physics and is not equivalent to the other two approaches. A more detailed discussion can be found in [1, 5]. Therefore, the relation between the LQS parameter  $\lambda_{LQS}$  [9] and the gluon density such a procedure would yield [1, 5] should be interpreted with great care. The connection between LQS and BDMPS/color dipole approach certainly needs further study.

## ACKNOWLEDGMENTS

This work was supported by the U.S. Department of Energy at Los Alamos National Laboratory under Contract No. W-7405-ENG-38.

## References

- [1] R. Baier, Y. L. Dokshitzer, A. H. Mueller, S. Peigne and D. Schiff, Nucl. Phys. B **484** (1997) 265 [arXiv:hep-ph/9608322].
- [2] F. Arleo, these proceedings (and references therein).
- [3] B. G. Zakharov, JETP Lett. **65** (1997) 615 [arXiv:hep-ph/9704255].
- [4] R. Baier, Y. L. Dokshitzer, A. H. Mueller and D. Schiff, Nucl. Phys. B **531** (1998) 403 [arXiv:hep-ph/9804212].
- [5] U. A. Wiedemann, Nucl. Phys. A **690** (2001) 731 [arXiv:hep-ph/0008241].
- [6] J. Dolejši, J. Hüfner and B. Z. Kopeliovich, Phys. Lett. B **312** (1993) 235 [arXiv:hep-ph/9305238].
- [7] M. B. Johnson, B. Z. Kopeliovich and A. V. Tarasov, Phys. Rev. C **63** (2001) 035203 [arXiv:hep-ph/0006326].
- [8] B. Blättel, G. Baym, L. L. Frankfurt and M. Strikman, Phys. Rev. Lett. **70** (1993) 896; L. Frankfurt, A. Radyushkin and M. Strikman, Phys. Rev. D **55** (1997) 98 [arXiv:hep-ph/9610274].
- [9] M. Luo, J. w. Qiu and G. Sterman, Phys. Rev. D **49** (1994) 4493.